

## **KYBELLA**

### **What is KYBELLA?**

KYBELLA is a prescription medicine used in adults to improve the appearance and profile of moderate to severe fat below the chin (submental fat), also called “double chin.”

### **Who should not receive KYBELLA?**

Do not receive KYBELLA if you have an infection in the treatment area.

### **Before receiving KYBELLA, tell your healthcare provider about all of your medical conditions, including if**

**you:** Have had or plan to have surgery on your face, neck, or chin; have had cosmetic treatments on your face, neck, or chin; have had or have medical conditions in or near the neck area; have had or have trouble swallowing; have bleeding problems; are pregnant or plan to become pregnant (it is not known if KYBELLA will harm your unborn baby); are breastfeeding or plan to breastfeed (it is not known if KYBELLA passes into your breast milk).

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take a medicine that prevents the clotting of your blood (antiplatelet or anticoagulant medicine).

### **What are the possible side effects of KYBELLA?**

**KYBELLA can cause serious side effects, including**

- Nerve injury in the jaw (which can cause an uneven smile or facial muscle weakness)
- Trouble swallowing
- Injection site problems including: bruising, hair loss, open sores (ulcers), damage and tissue cell-death (necrosis) around the injection site. Call your healthcare provider if you develop open sores or drainage from the treatment area

**The most common side effects of KYBELLA include:** Swelling, pain, numbness, redness, and areas of hardness in the treatment area. These are not all of the possible side effects of KYBELLA.

### **Procedure Time**

Depending on your specific targeted areas, a Kybella treatment usually takes about 45 minutes.

### **Side Effects & Recovery Times**

As with any other injection, clients may experience some side effects following their procedure. The most common side effects include mild to moderate redness, swelling, bruising, or numbness around the targeted injection area.

### **Results**

Typically, clients notice enhanced definition and contour in their chin’s profile after 2 to 4 treatments. Your injections will be spaced a minimum of 4 weeks apart and up to 6 treatments may be performed.